

Appendix A
Michigan Critical Health Indicators Summary Table

Indicator			Year	Rate/Percent/ Number
Focused Indicators	Health Risk Behaviors	Induced Abortions	1999	11.5/1,000
		Adolescent Use of Alcohol, Tobacco, Other Drugs	1999	
		Current drinker		49%
		Current smoker		34%
		Current marijuana user		26%
		Current cocaine user		3%
		Current Cigarette Smoking	2000	24.0%
		Overweight	2000	61.1
	Teen Pregnancy	1998	71.6/1,000	
	Morbidity & Mortality	AIDS Deaths*	1998	2.5/100,000
		Alcohol Induced Deaths*	1998	5.6/100,000
		Chlamydia Cases	1999	235.4/100,000
		Infant Mortality	1998	8.2/1,000
		Suicides*	1998	9.1/100,000
		Tuberculosis Cases	1999	3.6/100,000
	Preventive Health Care	Adequate Prenatal Care	1998	74.8%
		Childhood Immunizations	1999	
		4-3-1 Series		76%
		4-3-1-3-3 Series		71%
		Mammography (Never had a mammogram)	2000	8.8%
		Medicaid Managed Care Quality Assurance	1997	
		Well Child office visit		83%
		Up-to-date immunizations		79%
Otitis Media follow-up			67%	
MiChild Enrollment	2000	18,556		
Healthy Kids Enrollment	2000	274,164		
Vital Statistics Indicators	Heart Disease Deaths	1998	137.9/100,000	
	Cancer Deaths	1998	125.3/100,000	
	Stroke Deaths	1998	25.9/100,000	
	Chronic Obstructive Pulmonary Disease Deaths	1998	20.7/100,000	
	Unintentional Injury Deaths	1998	26.7/100,000	
	Pneumonia and Influenza Deaths	1998	13.1/100,000	
	Diabetes Related Deaths	1998	43.7/100,000	
	Kidney Disease Deaths	1998	5.0/100,000	
	Chronic Liver Disease and Cirrhosis Deaths	1998	8.0/100,000	
	Homicides	1998	8.4/100,000	
	* Mortality rates are age-adjusted, excluding infant mortality (See <i>Technical Notes</i> for definition).			

Last Updated: May 2002.